
Topical Anesthetic Cream

# MUCHCAINE







# Topical Anesthetic Cream





500g



30g

## Product Details

#### Composition


1 g contains

Active ingredient: Lidocaine — 10.56 mg
Additives (preservatives): Benzalkonium Chloride — 0.2 mg

#### Appearance

White or Pale Yellow Cream

## What Is a Lidocaine Cream?



Topical anesthetic molecule

A lidocaine cream is a topical anesthetic. It causes loss of feeling in the skin and surrounding tissues. It works by preventing nerves from transmitting painful impulses to the brain. It is used to reduce pain during and after some procedures.

#### Efficacy Effectiveness

- Topical anesthetic cream for needle (injection, blood test), minor skin operation, laser, waxing, tattoo, cosmetic procedures
- Cosmetic procedures, microdermabrasion, cosmetic procedures involving
- Medical procedures such as removal warts needles, microneedle therapy



500g

## Product Details

#### Composition

1 g contains

Active ingredient: Lidocaine \_\_\_\_\_\_\_ 25 mg

Active ingredient: Prilocaine \_\_\_\_\_\_\_ 25 mg

Additives (preservatives): Methylparaben \_\_\_\_\_\_\_ 1.6 mg

Additives (preservatives): Propylparaben \_\_\_\_\_\_\_ 0.4 mg

#### **Application**







Urology



Plastic Surgery



Tattoo

The effect of a lidocaine cream remains on the surface of the skin, so it is not used for pain that is felt under the skin. It is used to reduce pain during some procedures and treatments such as tattoos, laser/thread procedures, plastic surgery (injecting fillers), or ejaculation.

#### **Usage**

Apply before injections, or laser surgeries, or as directed by physicians. Results are obtained within 15 minutes, however, they may vary with factors such as skin color/sensitivity/thickness and acceptance to the active ingredient. Some procedures such as hair removal or tattoos may take more time once applied (15 minutes to 1 hour). No dressing is required. The maximum recommended dose for adults is 0.5 g.



#### 1. Do not use if:

- You are allergic (hypersensitive) to lidocaine or any other ingredient of the product
- There is skin inflammation
- You are pregnant

#### 2. Take special care if:

- You are a geriatric or pediatric patient
- You are a patient sensitive or allergic to propylene glycol

#### 3. General caution

- If any side effects bother you, please, stop using the cream and visit a doctor immediately
- Prolonged usage for more than 3 months without a consultation with a doctor is prohibited

#### 4. Application caution

- External use only. Do not apply on your eyes or nose. In case this happens, rinse thoroughly with water
- Do not use in large quantities, or on blistered areas









- Store sealed at 1-30°C
- Do not discard into fire
- Keep out of reach of children

